

[ET-36] The use of machine learning for auditing and modeling of vital sign data during open thoracotomy versus thoracoscopic surgical resections for congenital cystic lung lesions

<sup>1</sup>Simpao A, <sup>2</sup>Ivanov R, <sup>1</sup>Galvez J, <sup>2</sup>Lee I, <sup>1</sup>Laje P, <sup>1</sup>Flake A, <sup>3</sup>England W, <sup>3</sup>Choudhry D, <sup>3</sup>Wartman E, <sup>1</sup>Rehman M  
<sup>1</sup>Perelman School of Medicine at the Children's Hospital of Philadelphia and the University of Pennsylvania , Philadelphia , PA, United states; <sup>2</sup>University of Pennsylvania , Philadelphia , PA, USA; <sup>3</sup>The Children's Hospital of Philadelphia , Philadelphia , PA, United states

---

**Background:** Congenital cystic lung lesions warrant surgical resection to avoid the development of complications later in life (e.g. infection, pneumothorax, hemothorax, malignancy).[1] Thoracoscopic approach generally requires one-lung ventilation to optimize lung visualization.[2] Intraoperative loss of lung isolation is often unanticipated and usually results in conversion to open thoracotomy.

**Methods:** The anesthesia information management system data from patients with congenital lung lesions who underwent elective open and thoracoscopic lung lobectomy surgery between March 2005 and January 2012 at our tertiary care pediatric hospital were reviewed retrospectively. Existing implementations[3] of machine learning algorithms (logistic regression, support vector machine, perceptron) were used to predict the type of lobectomy (open versus thoracoscopic) via vital sign data patterns (Fig 1 and 2). The features chosen included the minimum SpO<sub>2</sub> value, the SpO<sub>2</sub> value 1, 2, 3, and 4 minutes after the minimum SpO<sub>2</sub> value, number of measurements less than 100% (normalized by operation length), and the mean and variance. The cases that were incorrectly identified by the learning algorithms were analyzed for possible causes.

**Results:** The data set consisted of 197 open thoracotomies and 95 thoracoscopic elective lobectomy cases. Prediction accuracy of thoracotomy versus thoracoscopic procedures was 80%. The algorithm misidentified more than half of the thoracoscopic cases as open cases, mainly due to minimal SpO<sub>2</sub> variation. Complications included thoracoscopic cases that were converted to open thoracotomies. During the data analysis, a few cases that were classified by the surgeon as strictly thoracoscopic were identified as open by the machine learning algorithms; these cases were confirmed via manual chart review as open cases.


**Conclusions:** While the use of SpO<sub>2</sub> as a feature of the classifier algorithm resulted in a fair accuracy rate, attempts should be made to improve the accuracy using other variables (e.g. EtCO<sub>2</sub>, respiratory rate, weight, age, tidal volume/weight). Future uses for the predictive algorithm include development of a smart alarm to warn providers of impending loss of lung isolation and one-lung ventilation, as well as the converse (e.g. mainstem intubation in an intubated patient). Future analysis should also lend insight into practices for optimal intraoperative management of one-lung ventilation in infants undergoing lung resection in order to decrease intraoperative SpO<sub>2</sub> variation and maximize oxygenation.

#### References

1. Parikh D, Samuel M. Congenital cystic lung lesions: Is surgical resection essential? *Pediatr Pulmonol.* 2005;40:533-7.
2. Hammer GB. Single-lung ventilation in infants and children. *Paediatr Anaesth.* 2004;14:98-102.
3. Mark Hall, et al. The WEKA Data Mining Software: An Update. *SIGKDD Explorations.* 2009;11:1.

# SpO2 and EtCO2 Patterns

## Thoracoscopic



## Open thoracotomy



SpO2 – blue  
EtCO2 - red